

## University of Groningen

### Birth and death of cellular senescence

Wang, Boshi

DOI:  
[10.33612/diss.131223530](https://doi.org/10.33612/diss.131223530)

**IMPORTANT NOTE:** You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

*Document Version*  
Publisher's PDF, also known as Version of record

*Publication date:*  
2020

[Link to publication in University of Groningen/UMCG research database](#)

*Citation for published version (APA):*  
Wang, B. (2020). *Birth and death of cellular senescence*. [Thesis fully internal (DIV), University of Groningen]. University of Groningen. <https://doi.org/10.33612/diss.131223530>

#### Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

#### Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

---

## CHAPTER II

### **Senescent cells in cancer therapy: friends or foes?**

*Boshi Wang<sup>#</sup>, Jaskaren Kohli<sup>#</sup> and Marco Demaria<sup>\*</sup>*

*European Research Institute for the Biology of Ageing (ERIBA), University Medical Center  
Groningen, 9713AV Groningen, The Netherlands*

*<sup>#</sup>Equal contribution*

*<sup>\*</sup>Correspondence to: [m.demaria@umcg.nl](mailto:m.demaria@umcg.nl)*

---

*Published in <Trends in Cancer> (2020)*

*(DOI:<https://doi.org/10.1016/j.trecan.2020.05.004>)*

- **Highlights**

- Cellular senescence is a common outcome of various anticancer interventions;
- Senescence-associated secretory phenotypes (SASPs) have pro-tumorigenic functions;
- Evidence exists of increased cellular senescence in patients treated for various types of cancer;
- Therapy-induced senescence can cause cancer metastasis and relapse and several adverse reactions to cancer treatments;
- Pharmacological interference with detrimental senescence might be considered to improve the efficacy of cancer treatments and improve the life quality of treated patients.

- **Abstract**

Several cancer interventions induce DNA damage and promote senescence in cancer and non-malignant cells. Senescent cells secrete a collection of pro-inflammatory factors collectively termed the Senescence-Associated Secretory Phenotype (SASP). SASP factors are able to potentiate various aspects of tumorigenesis, including proliferation, metastasis and immunosuppression. Moreover, accumulation and persistence of therapy-induced senescent cells can promote tissue dysfunction and early onset of various age-related symptoms in treated cancer patients. Here, we review in detail the mechanisms by which cellular senescence contributes to cancer development and side effects of cancer therapies. We also review how pharmacological interventions to eliminate senescent cells or inhibit SASP production can mitigate these negative effects, and propose therapeutic strategies based on the age of the patient.

Key words: cellular senescence, SASP, cancer therapy, pro-tumorigenesis, senotherapy

- **Cellular senescence**

Cellular senescence is a multifaceted and highly heterogeneous state characterized by a generally irreversible growth arrest, elevated lysosomal activity, resistance to apoptotic stimuli, deregulated metabolism, persistent DNA damage and elevated secretion of chemokines, cytokines and growth factors [1]. The stability of the cell cycle arrest relies on the CDK inhibitors p16 and p21 - often regulated by the master tumor suppressor protein p53 – and their elevated expressions serve as markers for senescence detection [2]. Another important feature of virtually all senescent cells is the Senescence-Associated (SA)- $\beta$ -galactosidase, a lysosomal enzyme whose activity is increased in senescent cells. Especially in cancer studies, SA- $\beta$ -galactosidase activity is often used as a unique surrogate to define a senescence state. However, SA- $\beta$ -galactosidase activity can be observed in some non-senescent contexts such as hair follicles and sebaceous glands in the skin [3]. It can also be detected in activated macrophages including those infiltrating tumors, which may provide difficulties in precise identification of senescent cancer cells in a tumor microenvironment *in vivo* [4,5]. Recent studies have demonstrated that senescent cells are able to resist pro-apoptotic stresses via the upregulation of anti-apoptotic mechanisms. Specific regulators of this function seem highly cell type- and stress-dependent, making their use as markers for senescence challenging [6,7]. On the same line, while metabolism is perturbed in senescent cells, no clear common metabolic mechanism that can be exploited for senescence identification has been demonstrated. A common trigger

for senescence induction is the activation of a persistent DNA damage response (DDR), making the DDR markers  $\gamma$ -H2AX or 53BP1 foci staining widely used markers for senescent cells. These proteins are also components of the DNA-SCARS (DNA Segments with Chromatin Alterations Reinforcing Senescence) [8]. Moreover, senescence-associated heterochromatin foci (SAHF), which are regions of condensed chromatin containing heterochromatin protein 1 and the histone H2A variant macroH2A and HMGA proteins, are observed in human, but not mouse, senescent cells [9]. One of the most variable senescence-associated phenotypes is the SASP, a transcriptional program for genes encoding secreted proteins which is a consequence of persistent DDR [10], and mediated by the NF- $\kappa$ B [11], p38 MAPK [12] and C/EBP $\beta$  pathways [13]. The SASP is enriched in pro-inflammatory factors, such as IL6, IL8, CXCL1, CCL2, CCL5 and MMP3 [14], which can be used as markers for DNA damage-induced senescence. The senescence-associated growth arrest is seen as a potent strategy to interfere with cancer progression. Moreover, SASP factors can stimulate immunosurveillance mechanisms and potentiate the tumor suppressive function of senescent cells by guiding the immune system to clear cancer cells. For this reason, features of senescent cells serve as common biomarkers to evaluate the efficacy of anti-cancer treatments. While cellular senescence is an oncosuppressive mechanism and a potent anti-cancer therapeutic strategy (for more information see [15,16]), several issues exist with generation and persistence of therapy-induced senescent cells. First, the SASP can play detrimental roles, and protect tumors from immune clearance, provide growth factors and enhance angiogenesis. Second, because most anti-cancer treatments are administered via systemic routes, many senescent cells are generated in non-tumor areas [17]. This excessive accumulation can have substantial side effects and accelerate the onset and progression of chronic diseases, such as cardiovascular diseases and neurodegenerative diseases, which normally are observed at advanced age [18]. Third, because cancer cells are genomically unstable, they can bypass the senescence growth arrest and restore aggressive and uncontrolled proliferation. This review will analyze these points and discuss the potential strategies to limit the detrimental effects of therapy-induced senescent cells.

Table 1, Summary of senescence-inducing cancer therapy

Category	Drug	Cell type	Markers	Treatments	Reference
<b>Chemotherapy</b>	Doxorubicin	Fibrosarcoma (HT1080)	(1) Elevated SA- $\beta$ -galactosidase activities (2) G1 cell cycle arrest (3) Increased micronuclei formation	40 nM for 4 days	[37]
		Breast cancer (MCF7)	(1) Elevated SA- $\beta$ -galactosidase activities (2) Increased micronuclei formation	50 nM for 2 days	

Breast cancer (MDA-MB-231)	(1) Increased micronuclei formation	50 nM for 2 days	
Colon cancer (HCT116)	(1) Elevated SA- $\beta$ -galactosidase activities (2) G1 cell cycle arrest (3) Increased micronuclei formation	50 nM for 4 days	
Prostate cancer (PC3)	(1) G1 cell cycle arrest (2) Increased micronuclei formation	100 nM for 4 days	
Prostate cancer (LNCaP)	(1) Elevated SA- $\beta$ -galactosidase activities (2) G1 cell cycle arrest (3) Increased micronuclei formation	100-500 nM for 4 days	
Ovarian cancer (A2780)	(1) Elevated SA- $\beta$ -galactosidase activities (2) G1 cell cycle arrest (3) Increased micronuclei formation	20-50 nM for 4 days	
Glioma (U251)	(1) Elevated SA- $\beta$ -galactosidase activities (2) Increased micronuclei formation	200 nM for 4 days	
Leukemia (K562)	(1) Elevated SA- $\beta$ -galactosidase activities	50 nM for 4 days	[21]
Melanoma (SK-MEL-103)	(1) Elevated SA- $\beta$ -galactosidase activities	1 $\mu$ M for 7 days	[31]
Vascular smooth muscle cells	(1) Elevated SA- $\beta$ -galactosidase activities (2) Morphology (enlarged) (3) G1 cell cycle arrest (4) p53/p21 upregulation (5) p16 upregulation (6) SASP induction (7) Increased DNA damage response	100 nM for 24 hours	[134]
Prostate stromal cells (PSC27)	(1) Elevated SA- $\beta$ -galactosidase activities (2) Reduced BrdU incorporation	10 $\mu$ M for 24 hours	[135]
Rat alveolar epithelial cells (L2)	(1) Elevated SA- $\beta$ -galactosidase activities (2) p53/p21 upregulation	50 nM for 24 hours	[136]
Human fibroblasts (IMR90)	(1) Elevated SA- $\beta$ -galactosidase activities (2) p16 upregulation (3) SASP induction	100 nM for 2 times 24 hours 24 hours drug holiday in between	[137]
Human fibroblasts (WI38)	(1) Elevated SA- $\beta$ -galactosidase activities (2) p53 upregulation	100 or 500 ng/ml for 4 days	[138]
Mouse dermal fibroblast	(1) Elevated SA- $\beta$ -galactosidase activities (2) Reduced EdU incorporation (3) p21 upregulation (4) p16 upregulation (5) <i>LMNB1</i> reduction (6) SASP induction (7) DNA damage response markers	250 nM for 24 hours	[29]
Reporter p16-3MR mice	(1) p16 upregulation in tissues (2) Increased bioluminescence (3) DNA damage response markers (4) SASP in serum (5) p21 staining in the heart and skin	10 mg/kg once	

Etoposide	Fibrosarcoma (HT1080)	(1) Elevated SA- $\beta$ -galactosidase activities (2) G1 cell cycle arrest (3) Increased micronuclei formation	900 nM Once	[37]
	Fibroblasts (IMR90)	(1) Elevated SA- $\beta$ -galactosidase activities (2) p53 upregulation	100 $\mu$ M Once	[11]
Camptothecin	Colon cancer (HCT116)	(1) Elevated SA- $\beta$ -galactosidase activities (2) Reduced colony formation capacity (3) p53/p21 upregulation	20 nM for 96 hours	[139]
	Lung cancer (H1299)	(1) Elevated SA- $\beta$ -galactosidase activities	30 nM for 24 hours or 48 hours	[22]
Cisplatin	Nasopharyngeal carcinoma (CNE1)	(1) Elevated SA- $\beta$ -galactosidase activities	0.5 or 2 $\mu$ g/ml for 24 or 48 hours	[20]
	Fibrosarcoma (HT1080)	(1) Elevated SA- $\beta$ -galactosidase activities (2) G1 cell cycle arrest (3) Increased micronuclei formation	2.2 $\mu$ M	[37]
	Immortalized skin fibroblasts (hTERT-BJ)	(1) Elevated SA- $\beta$ -galactosidase activities (2) p53 upregulation (3) IL6 upregulation	100 $\mu$ M for 24 to 72 hours	[140]
	Reporter p16-3MR mice	(1) Increased bioluminescence (2) p16 upregulation in skin and lung	2.3 mg/kg for 3 consecutive days	[29]
	Breast cancer (MCF7)	(1) Reduced proliferation (2) Increased DNA damage response (3) Elevated SA- $\beta$ -galactosidase activities	0.03, 0.3, 1 $\mu$ M for 5 days	[28]
Paclitaxel	Endothelial cell (HUVEC)	(1) Elevated SA- $\beta$ -galactosidase activities	2.5 or 5 nM for 24 hours	[23]
	Immortalized skin fibroblasts (hTERT-BJ)	(1) Elevated SA- $\beta$ -galactosidase activities (2) p53 upregulation (3) IL6 upregulation	100 nM for 24 to 72 hours	[140]
	Human bone marrow mononuclear cells	(1) Reduced proliferation (2) Increased DNA damage response (3) Elevated SA- $\beta$ -galactosidase activities	0.03, 0.3, 1 $\mu$ M for 5 days	[28]
	Mouse dermal fibroblast	(1) Elevated SA- $\beta$ -galactosidase activities (2) Reduced population doubling (3) p16 upregulation (4) <i>LMNB1</i> reduction (5) SASP induction (6) Increased DNA damage response	50 nM for 24 hours or 48 hours	[29]
	Reporter p16-3MR mice	(1) Increased bioluminescence (2) p16 upregulation in skin and lung	10 mg/kg for 3 consecutive days	
	Lung cancer (A549)	(1) Elevated SA- $\beta$ -galactosidase activities (2) Reduced BrdU incorporation (3) Increased cell size (4) p53/p21 upregulation	50 $\mu$ g/ml for 120 hours 50 mU/ml for 48 hours 0.05-50 $\mu$ g/ml for 72 hours	[32,141,142]
	Mouse lung tissue	(1) Elevated SA- $\beta$ -galactosidase activities (2) Upregulated p21 staining (3) Reduced p-RB staining (4) Increased DNA damage response	5 U/kg 50 mg/kg 2.5 mg/kg	[32–34]

		(5) Increased SASP		
Temozolomide	Rat alveolar epithelial cell / lung tissue	(1) Elevated SA- $\beta$ -galactosidase activities (2) Reduced BrdU incorporation	5 $\mu$ g/ml for 24 hours	[136]
	Human alveolar epithelial cell	(1) Elevated SA- $\beta$ -galactosidase activities (2) p21 upregulation (3) p16 upregulation	0.1-10 $\mu$ g/ml for 5 days	[32,142]
	Skin fibroblasts (HCA2)	(1) SASP induction	10 $\mu$ g/ml for 24 hours	[84]
	Prostate stromal cells (PSC27)	(1) Elevated SA- $\beta$ -galactosidase activities (2) Reduced BrdU incorporation (3) SASP induction	50 $\mu$ g/ml for 24 hours	[135]
	Glioma	(1) Elevated SA- $\beta$ -galactosidase activities (2) Reduced colony formation capacity (3) p53/p21 upregulation	10 $\mu$ g/ml for 24 hours 100 $\mu$ M for 24 hours	[143–146]
	Melanoma	(1) Reduced colony formation capacity (2) p53/p21 upregulation	25 to 250 $\mu$ M for 12 days	[147]
	Colorectal cancer	(1) Elevated SA- $\beta$ -galactosidase activities	100 to 1000 $\mu$ M for 48 hours	[24]
	Reporter p16-3MR mice	(1) Increased bioluminescence (2) p16 upregulation in skin and lung	10 mg/kg for 3 consecutive days	[29]
Radiotherapy	Fibrosarcoma (HT1080)	(1) Elevated SA- $\beta$ -galactosidase activities (2) G1 cell cycle arrest (3) Increased micronuclei formation	1300 rad	[37]
	Breast cancer (MCF7)	(1) Elevated SA- $\beta$ -galactosidase activities	5 times 2 Gy	[148]
	Breast cancer (MCF7 and MDA-231)	(1) Elevated SA- $\beta$ -galactosidase activities (2) p53 and p21 upregulation	10 Gy	[39]
	Glioblastoma (U87)	(1) Elevated SA- $\beta$ -galactosidase activities	2 Gy	[40]
	Endothelial cells (HUVEC and BAEC)	(1) Elevated SA- $\beta$ -galactosidase activities (2) Reduced BrdU incorporation	8 Gy	[41]
	Keloid fibroblasts	(1) Enlarged cell size (2) Elevated SA- $\beta$ -galactosidase activities (3) G1 cell cycle arrest (4) p16, p21 upregulation	2, 4, 8 Gy	[42]
	Fibroblasts (HCA2)	(1) Elevated SA- $\beta$ -galactosidase activities (2) SASP induction (3) Colony formation assay (irreversible)	10 Gy	[43]
	Fibroblasts (IMR90 and WI38)	(1) Elevated SA- $\beta$ -galactosidase activities (2) Reduced BrdU incorporation	2.5, 5, 7.5 and 10 Gy	[44]
	Mouse inguinal fat	(1) p16 upregulation in organs (2) Elevated SA- $\beta$ -galactosidase activities	10 Gy local irradiation on one leg	[45]
	Fibroblasts from naked mole rats	(1) Elevated SA- $\beta$ -galactosidase activities (2) p21 upregulation (3) Reduced BrdU incorporation	10 Gy or 20 Gy	[46]
	Mouse tibialis anterior muscle	(1) Elevated SA- $\beta$ -galactosidase activities	10 Gy local irradiation	[47]
	Reporter p16-3MR mice	(1) Increased bioluminescence (2) p16 upregulation in organs (3) SASP induction in organs	7 Gy total body irradiation	[48]



<b>CDK4/6 inhibitors</b>	Palbociclib	Mouse bone	(1) p16 upregulation in organs (2) SASP induction in organs	20 Gy local irradiation	[113]
		Breast cancer (MCF7)	(1) Elevated SA- $\beta$ -galactosidase activities (2) Reduced Proliferation	0.1, 0.3, 1 $\mu$ M for 5 days	[28]
		Breast cancer (MCF7 and T47D)	(1) Elevated SA- $\beta$ -galactosidase activities (2) Reduced Proliferation (reversible)	1 or 5 $\mu$ M for 6 days *5 $\mu$ M shows off-target effects	[149]
		Melanoma (MEL10)	(1) Elevated SA- $\beta$ -galactosidase activities (2) Up-regulated cyclin-D1	0.5 $\mu$ M for 7 days	[150]
		Melanoma (1205Lu)	(1) G1 growth arrest (FACS) (2) Reduced p-Rb (3) Elevated SA- $\beta$ -galactosidase activities (4) SASP induction (5) Did not check reversibility	1 $\mu$ M for 8 days	[68]
		Melanoma (SK-MEL-103)	(1) Elevated SA- $\beta$ -galactosidase activities	1 $\mu$ M for 14 days	[31]
		Melanoma (SK-MEL-103)	(1) Elevated SA- $\beta$ -galactosidase activities (2) Colony formation assay	1 $\mu$ M for 7 days	[21]
		Liposarcoma (LS8817, LS141, LS0082)	(1) Elevated SA- $\beta$ -galactosidase activities (2) Reduced BrdU incorporation	1 $\mu$ M for 2 days	[151]
		Liposarcoma (U2OS)	(1) Elevated SA- $\beta$ -galactosidase activities	1 $\mu$ M for 7 days	[152]
		Liposarcoma (LS8107)	(1) Elevated SA- $\beta$ -galactosidase activities	1 $\mu$ M for 7 days	[153]
		Gastric cancer (AGS and MKN-45)	(1) Elevated SA- $\beta$ -galactosidase activities (2) Reduced proliferation	0.5 $\mu$ M for 2 or 4 days	[154]
		Hepatocellular carcinoma (Huh7 and skHep1)	(1) Elevated SA- $\beta$ -galactosidase activities (2) Reduced proliferation (drug on) (3) Colony formation assay (drug off) (Irreversible) (4) Cyclin D1 upregulation	0.1, 0.5, 1 $\mu$ M for 10 or 14 days	[155]
		Human bone marrow mononuclear cells (hBMNC)	(1) Reduced proliferation (reversible) (2) NO DNA damage response (3) NO SA- $\beta$ -galactosidase activities	0.1, 0.3, 1 $\mu$ M for 5 days	[28]
		Mouse embryonic fibroblast	(1) Elevated SA- $\beta$ -galactosidase activities (2) G1 cell cycle arrest (drug on) (3) NO DNA damage response (4) p16 upregulation (5) p21 upregulation (6) SASP induction	4 $\mu$ M for 2 times 4 days	[64]
		Human fibroblasts (IMR90)	(1) NO SASP induction detected	10 $\mu$ M for 7 days	[65]
	Ribociclib	Neuroblastoma	(1) Elevated SA- $\beta$ -galactosidase activities	500 nM for 6 days	[54]
		Ewing Sarcoma (SKNEP1)	(1) Elevated SA- $\beta$ -galactosidase activities (2) Cyclin D3 upregulation	1 $\mu$ M for 5 days	[58]
	Abemaciclib	Breast cancer (MDA-MB-453)	(1) Elevated SA- $\beta$ -galactosidase activities	25 nM for 48 hours	[59]
		Breast cancer xenograft (MDA-MB-453)	(1) Elevated SA- $\beta$ -galactosidase activities (2) Reduced KI67 <sup>+</sup> cells	90 mg/kg daily for 5 days	[60]

<b>Epigenetic modulator</b>	5-aza-2'-deoxycytidine	Breast cancer (MDA-MB-453 and BT474)	(1) Elevated SA- $\beta$ -galactosidase activities (2) Reduced proliferation (reversible)	250 or 500 nM for 7 days	[60]
		Breast cancer transgenic model (MMTV-rtTA/tetO-HER2)	(1) Elevated SA- $\beta$ -galactosidase activities (2) NO SASP induction detected	75-90 mg/kg daily for 12 days	[60]
		Breast cancer (MDA-MB-453)	(1) Elevated SA- $\beta$ -galactosidase activities	500 nM for 72 hours	[61]
		Breast cancer (MCF7)	(1) Elevated SA- $\beta$ -galactosidase activities	500 nM for 4 doubling time	[62]
		Breast cancer (MCF7)	(1) G1 cell cycle arrest (2) NO SASP induction detected	500 nM for 6 hours	[156]
	Suberoylanilide hydroxamic acid (SAHA)	Osteosarcoma (U2OS)	(1) Elevated SA- $\beta$ -galactosidase activities (2) p16 upregulation	5 $\mu$ M or 10 $\mu$ M	[69]
		Lung mesothelioma (H28)	(1) Elevated SA- $\beta$ -galactosidase activities (2) p21/p27 upregulation (3) Reduced proliferation (4) DNA damage response	1 $\mu$ M or 10 $\mu$ M for 2 days	[70]
		Liver cancer (HepG2, Hep3B and Huh-7)	(1) Elevated SA- $\beta$ -galactosidase activities (2) SASP induction (3) Reduced proliferation	20 $\mu$ M for 24 hours 12 $\mu$ M for 72 hours	[157,158]
		Colon cancer (HCT116)	(1) Elevated SA- $\beta$ -galactosidase activities (2) Reduced proliferation (3) Colony formation assay	2 $\mu$ M for 24 hours	[73]
		Leukemia (MOLM-7, HL-60 and JURL-MK1)	(1) Reduced proliferation (2) Elevated SA- $\beta$ -galactosidase activities	0.5-10 $\mu$ M for 72 hours	[74]
		Urothelial carcinoma	(1) Re-plated colony formation assay (2) G1 cell cycle arrest	2.5 $\mu$ M for 48 hours	[75]
		Glioblastoma stem cell	(1) Elevated SA- $\beta$ -galactosidase activities	2.5 $\mu$ M for 7 days	[76]
		Adenoid cystic carcinoma primary cell	(1) Elevated SA- $\beta$ -galactosidase activities (2) p16 upregulation (3) DNA damage response	1 or 1.4 $\mu$ M for 120 hours	[72]
<b>Immunotherapy</b>	Rituximab (CD20-targeting antibody)	Human lymphoma (EHEB, RC-K8, SD1)	(1) Elevated SA- $\beta$ -galactosidase activities (2) DNA damage responses (3) SASP induction (4) p53 and p21 induction	20 $\mu$ g/ml	[79]

## • Cancer interventions and induction of senescence

A significant number of commonly used cancer interventions has been associated to induction of cellular senescence in either tumor or non-tumor cells and tissues (Table 1). These senescence-inducing interventions can be categorized in 5 families: chemotherapy, radiotherapy, CDK4/6 inhibitors, epigenetic modulators, and immunotherapy.

### 1) Chemotherapy

Chemotherapy is defined as the use of cytotoxic compounds to inflict unrepairable DNA damage or impair mitosis in cancer cells. It can be originated to the 1940s when nitrogen mustard was used to treat non-Hodgkin's lymphoma; later work found this was due to its ability to alkylate DNA (alkylation is described in more detail below) [19]. Over time, other types of cytotoxic agents have been developed and adopted. Numerous studies demonstrated that chemotherapeutics can induce senescence in both malignant and non-malignant cells, but in many reports the only evidence of senescence is elevated SA- $\beta$ -galactosidase activity [20–24]. Few analyses also measured the level of SASP expression, a phenotype which could provide information on the non-autonomous effect of senescent cancer and non-cancer cells.

Topoisomerase inhibitors block topoisomerase enzymes from re-ligating DNA strands after supercoil unwinding, thereby preventing replication [25]. Non-sealed DNA is recognized as a break and activates DDR signaling. Doxorubicin is the most common topoisomerase inhibitor and is used to treat a variety of cancers including lung, breast, lymphoma and acute lymphocytic leukemia [25]. Doxorubicin is capable of inducing senescence in various cancer cell types *in vitro* (Table 1). Additionally, doxorubicin is able to induce senescence in non-malignant cells such as vascular smooth muscle cells, prostate stromal cells, human and mouse dermal fibroblasts as well as tissues of cancer-free mice (Table 1). Etoposide and Camptothecin are topoisomerase inhibitors that are also used to treat a range of similar cancers to doxorubicin. These drugs have also been found to induce senescence of colon, breast, ovarian adenocarcinoma, hepatocarcinoma and osteosarcoma cells *in vitro*. Etoposide also induces senescence in primary fibroblasts (Table 1).

Alkylating agents are reactive compounds which form DNA cross-links by chemically reacting to atoms in DNA. Cross-linked DNA strands can break during DNA replication when the cell attempts division, resulting in DDR initiation [26]. Cisplatin and Temozolomide are typical alkylating agents that can induce cellular senescence (Table 1).

Paclitaxel belongs to a family of microtubule inhibitors used to treat patients with ovarian, breast, brain and non-small-cell lung cancers. The drug interferes with normal microtubule spindle dynamics which compromises the metaphase-anaphase transition, thereby arresting the cell at mitosis [27]. Treatment of MCF7 breast cancer cells [28] and various non-malignant cell lines such as endothelial cells, bone marrow mononuclear cells, as well as human and mouse fibroblasts induces senescence [23,28,29]. Paclitaxel also increased bioluminescence in

p16 reporter mice, thereby concluding that the drug can induce senescence in non-cancerous tissues *in vivo* [29] (Table 1).

Bleomycin is commonly used to treat Hodgkin's lymphoma and testicular germ cell tumors. The drug oxidizes DNA nucleotides to induce single and double stranded breaks [30]. Bleomycin induces senescence in A549 lung cancer cells as shown by increased SA- $\beta$ -galactosidase and upregulation of p21. Bleomycin is also commonly used to induce pulmonary fibrosis in cancer-free mice [31] and induces senescence in human alveolar cell lines and mouse lung tissue. Importantly, as well as demonstrating senescence by SA- $\beta$ -galactosidase staining and p16 and p21 upregulation, some of these studies provided evidence of a SASP in lungs from treated mice [32–34] (Table 1). This is important considering lung damage and pulmonary inflammation are major side effects in human patients [35].

## 2) Radiotherapy

Radiotherapy entails the use of high energy electrically-charged ions to inflict irreparable DNA damage in cancer cells with the intent of inducing cell death. Because of its lack of specificity, radiotherapy can be used to treat numerous cancers including lymphoma, soft tissue sarcomas, central nervous system tumors and a wide range of carcinomas (lung, prostate, skin, breast, head and neck, bladder) [36]. As radiotherapy is locally administered to the site of the cancer, rather than systemically as with chemotherapy, it is less likely that non-cancerous tissue areas are damaged. Numerous studies have demonstrated that ionizing radiation induces SA- $\beta$ -galactosidase in breast, colon carcinoma, neuroblastoma and fibrosarcoma cell lines (Table 1) [37–39]. Interestingly, the senescence response to irradiation appears to be dependent on p53 status. Breast cancer cells carrying missense mutation in the p53 DNA binding domain undergo apoptosis instead of senescence [39], while glioblastoma cells with p53 mutation in the DNA binding domain recover from irradiation and resume proliferation [40]. It is possible that the cell fate decision in response to radiotherapy in p53 mutant tumors depends on the cancer type or on additional mutations affecting survival pathways. Future work should further characterize what variables influence the response to radiotherapy in order to select for treatment the patient population most likely to favorably respond to the treatment.

Ionizing radiation (IR) is commonly used to prematurely induce senescence in cells *in vitro* and *in vivo*. IR-treated cells show numerous markers of senescence including SA- $\beta$ -galactosidase, p16, p21 and SASP induction [29,41–47]. Cancer-free mice irradiated with sub-

lethal doses of  $\gamma$ -radiation also show several signs of senescence in various tissues, including p16 up-regulation [48–50] (Table 1).

### 3) CDK4/6 inhibitors

Cyclin-dependent kinases (CDK) 4 and 6 are critical factors for cell cycle transition from G1 to S phase [51]. Hyperactivation of the D type cyclin-CDK4/6 axis in cancer cells has been observed in many preclinical studies, particularly in the context of breast cancer [52,53], making its targeted inhibition an attractive anti-cancer strategy. Recently, three pharmacological inhibitors (palbociclib, ribociclib and abemaciclib) have been developed and approved by the U.S. FDA. Palbociclib was approved in combination with letrozole for estrogen receptor positive (ER<sup>+</sup>) advanced breast cancer patients in 2015. Two years later, ribociclib was approved in combination with an aromatase inhibitor to treat hormone receptor positive (HR<sup>+</sup>) human epidermal growth factor receptor 2 negative (HER2<sup>-</sup>) metastatic breast cancer patients. In the same year, the last CDK4/6 inhibitor abemaciclib got the approval in combination with fulvestrant for treating HR<sup>+</sup>;HER2<sup>-</sup> advanced breast cancer patients who failed to respond to endocrine therapy [54–56]. Because CDK4/6 inhibitors mimic the mechanism of action of p16 [57], induction of cellular senescence has been considered a likely outcome of the treatment. Indeed, palbociclib was shown to induce cellular senescence in breast and gastric cancer, melanoma, liposarcoma, and hepatocellular carcinoma, ribociclib in neuroblastoma cells [54] and Ewing Sarcoma cells [58] and abemaciclib in different breast cancer cells [59–62] (Table 1). However, senescence induction was mainly evaluated based on SA- $\beta$ -galactosidase positivity, and successful establishment of an irreversible proliferative arrest was rarely measured. While enhanced SA- $\beta$ -galactosidase activity remains one of the best senescence-associated features, its use as a sole and unequivocal marker for senescent cells remains debated [1,2]. For example, conditions of increased lysosomal content or activity might also lead to SA- $\beta$ -galactosidase activity. This has been demonstrated for CDK4/6 inhibitors in SK-Mel-103 melanoma cells and normal cells, where palbociclib and abemaciclib are trapped in the lysosomes and might induce lysosomal stress unrelated to growth arrest [63]. However, long-term treatments with palbociclib and abemaciclib are sufficient to induce senescence even when extra drug washouts to avoid lysosomal accumulation are applied [63]. Additional evidences for palbociclib able to promote a stable growth arrest derive from experiments using either mouse embryonic fibroblast [64] or IMR90 human fibroblasts [63,65]. These data are in-line with the notion that prolonged overexpression of p16 (for 7 days) is

needed to achieve permanent growth arrest, after which p16 itself becoming dispensable [66]. While the duration of the treatment is possibly an important determinant of cell fate, other variables might contribute to the sensitivity to CDK4/6 inhibitors. For example, palbociclib was also shown to induce reversible growth arrest in bone marrow mononuclear cells but irreversible growth arrest and senescence in MCF7 cells [28] (Table 1), suggesting a cell type-dependent mode of action. Studies on how senescence is induced by CDK4/6 inhibitors should be focusing on additional molecular and environmental conditions. Breast cancer patients carrying mutations in the coding region of p53 DNA-binding domain are poor responders to abemaciclib [67]. However, it was shown that palbociclib-induced senescence is independent from the p53 pathway, since prolonged treatment with palbociclib promoted G1 arrest in p53-deficient melanoma cells [68]. Nevertheless, the stability of the growth arrest after drug withdraw, which would suggest successful induction of senescence, was not verified.

#### 4) Epigenetic modulators

5-aza-2' deoxycytidine (5-aza) is an inhibitor of DNA methyltransferase (DNMT), which can cause global demethylation of CpG-enriched promoters or enhancers [69]. Since the p16<sup>Ink4a</sup> promoter contains CpG-rich regions, it was postulated that treatment with 5-aza could induce senescence by activating p16. Indeed, treatment with 5-aza was shown to induce p16 up-regulation and growth arrest in osteosarcoma cells (U2OS) [69] and elevated SA- $\beta$ -galactosidase activity and increased DNA damage response in lung mesothelioma cells (H28) [70]. Notably, the p53-p21 pathway was also activated by 5-aza treatment in U2OS and MCF7 cells that entered a senescence state [71] (Table 1). Suberoylanilide hydroxamic acid (SAHA), also known as vorinostat, is a histone deacetylase (HDAC) inhibitor that was approved by the U.S. FDA to treat cutaneous T cell lymphoma [72]. SAHA was shown to induce various senescence phenotypes in colon cancer cells (HCT116) [73], and senescence-like growth arrest in human leukemia cell lines (MOLM-7, HL-60 and JURL-MK1) [74] and urothelial carcinoma cells [75]. Interestingly, similar pro-senescence effects of SAHA were also observed for glioblastoma stem cells and adenoid cystic carcinoma primary cells [72,76] (Table 1). Other HDAC inhibitors, such as sodium butyrate and trichostatin-A, were initially tested in normal cells and shown to induce senescence in human normal fibroblasts [77,78].

#### 5) Immunotherapy

As immunotherapeutic drugs mediate immune-mediated cancer cell apoptosis, it is possible that the SASP could be a modulator of this process. Rituximab is a CD20 targeting antibody used to treat leukemia and lymphoma. Interestingly this drug has been shown to induce senescence in B cell lymphoma lines, as judged by increased SA- $\beta$ -galactosidase activity, p53/p21 signaling, DDR response and indeed a SASP [79](Table 1). Owing to these results, it would be warranted to investigate whether other immunotherapies induce senescence and SASP in other cancer types.

- **Pro-tumorigenic effects of senescent cells**

Inflammatory SASP factors secreted by senescent cells can promote various aspects of tumorigenesis in a non-autonomous manner, including cancer cell proliferation, migration, invasiveness, angiogenesis, epithelial-mesenchymal transition (EMT) as well as immune-mediated clearance.

The first evidence that the SASP can promote proliferation and growth of cancer cells came from studying co-cultures of preneoplastic epithelial cell lines with senescent WI38 lung fibroblasts. When cultured together *in vitro*, the proliferation rate of preneoplastic epithelial cells was quicker compared to co-cultures with non-senescent fibroblasts [80]. Importantly, this pro-growth effect was independent from the senescence-inducing stimulus ( $H_2O_2$ , RAS<sup>V12</sup> or p14<sup>ARF</sup> overexpression) but was not observed when senescent fibroblasts were co-cultured with normal epithelial cells [80]. In a xenograft murine model, faster growing epithelial tumors were observed when co-injected with senescent fibroblasts [80]. IL-6 is a key SASP factor which can promote proliferation in a paracrine manner, through binding to the IL-6 receptor and subsequent activation of STAT3 (Figure 1). STAT3 transcribes oncogenes and growth regulators including c-myc, c-Fos, cyclin D1 and mTORC1 [81]. Co-injection of breast cancer cells with senescent fibroblasts (induced by  $\gamma$ -irradiation or bleomycin) in nude mice also results in faster growing tumors [82].

The pro-tumorigenic SASP is also able to positively influence invasive properties of cancer cells. Breast cancer cells displayed an increased ability to migrate across a porous membrane when co-cultured with senescent cells [14]. These actions are reported to be modulated by various SASP factors. IL6 and IL8 seem to be major players, as invasiveness is rescued when blocking antibodies against these two interleukins are administered, while addition of recombinant IL6 or IL8 promotes invasiveness of preneoplastic epithelial cells co-cultured

with non-senescent fibroblasts [14]. IL-6 and IL-8 are able to mediate the activation of STAT3, which consequently drives transcription of various MMPs and promote cancer cell invasion [81,83]. For many decades MMPs have been known to positively influence cancer invasiveness through extracellular matrix degradation, allowing dissemination of cancer cells from the primary site [82]. It is also suggested that extracellular matrix degradation increases the permeability of adjacent capillaries, boosting access to mitogens and growth factors, such as HGF [84] (Figure 1).

The SASP also contributes to tumorigenesis by promoting EMT. Treating non-aggressive breast cancer cells with conditioned media from senescent fibroblasts induced various hallmarks of EMT including reduced  $\beta$ -catenin and E-cadherin and increased vimentin expression, consistent with cells acquiring a mesenchymal phenotype [14]. Interestingly, senescent cancer cells can also promote EMT in non-senescent counterparts. Incubating proliferating mesothelioma cells with conditioned media from mesothelioma cells induced to senescence by treatment with the chemotherapeutic pemetrexed results in gain of various EMT markers including vimentin upregulation [85]. Since STAT3 is a key transcriptional activator of EMT genes, IL-6 and IL-8 are therefore likely to be key SASP factors involved for this process. Indeed, IL-6 and IL-8 have been shown to induce EMT in cancer cells *in vitro* [86–88] (Figure 1).

Vascular endothelial growth factor (VEGF) is a factor which promotes angiogenesis and provides access to growth factors, eventually favoring malignant growth [89]. VEGF is also part of the SASP [14] and co-injection of senescent fibroblasts with malignant epithelial cells was shown to promote formation of blood vessels [90] (Figure 1).

- **Evidence of senescent cells in patients**


Considering the potential detrimental role of cellular senescence in promoting various aspects of tumorigenesis, more interest is raising on measuring senescence burst as a biomarker of toxicity. Prostate cancer patients treated with the chemotherapeutic mitoxantrone showed increased expression of various senescence-associated genes *p16*, *p21* and numerous SASP markers including *IL-6* and *IL-8* [14]. SA- $\beta$ -galactosidase,  $p16^+$  and  $p53^+$  cells have been detected in tumor tissues of breast cancer patients treated with cyclophosphamide, doxorubicin and 5-fluorouracil [16]. An increase in  $p16^+$ ,  $CD3^+$  T lymphocytes was also detected in breast cancer patients undergoing doxorubicin treatment [91]. Therefore, chemotherapy induces



senescence in both malignant and non-malignant cells in humans. *VEGF* and *CCL2* plasma concentrations in the breast cancer patients were significantly upregulated after treatment commenced in plasma. Importantly these markers of senescence remained consistent even 12 months after treatment finished [91], suggesting these senescent cells persist and are refractory to clearance.

SA- $\beta$ -galactosidase and p16 positive cells are located at the invasive border of untreated human papillary thyroid carcinoma tissues, where they are found to promote cancer cell metastasis [92]. It is unclear where these senescent cells originate from but since they are positive for *BRAF*<sup>V600E</sup> [92], they could be thyrocytes that have undergone oncogene-induced senescence (OIS). These cells may secrete pro-inflammatory factors which promote invasion and metastasis of local cancer cells that have bypassed OIS from gain of subsequent mutations in senescence maintenance genes.

Importantly, the presence of senescent cells in the tumor microenvironment is associated with worse disease-free survival. Human HCC patients had worse overall survival and shorter recurrence-free survival when their tumors displayed higher expression of genes involved in senescence. This was presumably due to the presence of an immunosuppressive environment [93].



**Figure 1. Pro-tumorigenic effects of senescent cells.** SASP factors are engaged in senescent cells through the activation of NF- $\kappa$ B, C/EBP $\beta$  and p38MAPK pathways. SASP factors contribute to different aspects of cancer progression: IL6 and IL8 promotes EMT; IL6-IL6R-STAT3 / CCL5-CCR5-c-Myc / HGF contribute to the proliferation of cancer cells; CXCL5 and VEGF promote angiogenesis; CXCL12-CXCR4 / IL6 / IL8 / Eataxin / CCL5 / MMPs promote the invasion and migration of cancer cells; IL6 suppresses the CD45<sup>+</sup>CD3<sup>+</sup> T cell-mediated immune clearance of cancer cells by activating the Gr-1<sup>+</sup> myeloid cells. CCL2 suppresses the NK cell-mediated immune clearance of cancer cells by activating the CCR2<sup>+</sup> myeloid cells. SASP is a collection of several factors which can collectively act. SASP, Senescence-Associated Secretory Phenotypes. MMP, matrix metalloproteinases. IL, interleukins. CCL, (C-C motif) chemokine ligands. CCR, (C-C motif) chemokine receptors. CXCL, (C-X-C motif) chemokine ligands. CXCR, (C-X-C motif) chemokine receptors. EMT, epithelial-mesenchymal transition.

- **Roles of senescent cells in side effects of cancer therapies**

### 1) Pro-tumorigenic functions

While the studies described in previous sections highlight the potential pro-tumorigenic function of senescent cells and the evidence of the presence of senescent cells in treated cancer patients, the use of more sophisticated *in vivo* systems has recently allowed us to test the pathological and clinical relevance of therapy-induced senescence.

In the breast carcinoma mouse model MMTV-Wnt1, doxorubicin induces senescence, as measured by elevated SA- $\beta$ -galactosidase activity, and p21 expression and SASP factors Eotaxin, CXCL5 and CCL5 expression. Interestingly, these SASP factors mediate cancer relapse by stimulating malignant transformation of surrounding non-senescent breast cancer cells [94] (Figure 1): Eotaxin promotes cancer cell invasion through a CCR3-ERK-mediated MMP3 activation [95]; CXCL5 stimulates angiogenesis through activation of VEGF [96] and metastasis via activation of AKT/GSK3 $\beta$ / $\beta$ -catenin signaling [97]; CCL5 promotes proliferation through upregulation of cyclin D1 and c-myc and invasion through increased expression of MMP-2 and MMP-9 [98]. Genetic and pharmacological clearance of doxorubicin-induced senescent non-malignant cells from an orthotopic model of breast carcinoma reduces cancer growth and limits cancer relapse [29]. Therefore, induction of senescence in non-malignant and malignant cells may promote tumorigenesis and cancer relapse via the SASP.

Loss-of-function mutations and deletions affecting *PTEN* are a common feature of human prostate cancer cells, accounting for up to 40% of primary tumors [99]. Homozygous inactivation of *PTEN* in mouse prostate epithelial cells promotes the formation of adenocarcinomas *in vivo* [100,101]. Interestingly, docetaxel, a chemotherapeutic agent commonly used for prostate cancer treatment, is unable to efficiently induce senescence in these PTEN-null tumors [100]. A major mechanism explaining this protection is the infiltration of Gr-1<sup>+</sup> myeloid cells which secrete interleukin receptor antagonist (IL-1RA) to block IL-1 $\alpha$ -mediated signaling and protect prostate cancer cells from establishment of senescence. Importantly, blocking myeloid cell recruitment via CXCR2 inhibition sensitizes prostate cancer cells to docetaxel-induced senescence [101].

Forced expression of *NRAS*<sup>G12V</sup> in hepatocytes led to senescence and secretion of CCL2, which, in turn, attracts CCR2<sup>+</sup> myeloid cells and promotes senescent cell clearance. However, these


immature myeloid cells promote growth of nearby hepatocellular carcinoma (HCC) cells by inhibiting NK cell-mediated cytotoxicity [93] (Figure 1). Myeloid infiltration was also found in a *KRAS*<sup>G12V</sup> lung adenoma model [101], suggesting this immunosuppressive mechanism could occur independently of type of cancer and/or oncogenic mutation.

p27 induction in mesenchymal cells promotes stromal senescence in skin and an immunosuppressive environment, due to recruitment of CD11b<sup>+</sup> and Gr-1<sup>+</sup> myeloid cells and decrease in CD45<sup>+</sup>CD3<sup>+</sup> lymphocytes. Myeloid recruitment is partly mediated by IL-6 secretion from senescent stromal cells (Figure 1). Co-injection of isolated senescent fibroblasts with skin carcinoma cells in mice results in a suppressive myeloid cell-mediated increase in tumor growth, and this effect was absent in immunocompromised mice [102]. These data suggest that senescence-inducing interventions might skew the immune system in a tumor-permissive mode, and warrant further investigation on the role of immune modulatory drugs to counteract the tumor promoting function of senescent cells.

IL-6 has also been reported to promote metastasis of breast cancer cells to bone sites [103]. Using a conditional mouse model where osteoblasts can be specifically induced to senescence through p27 expression, implanted breast cancer cells metastasized at greater frequencies in the bone when p27 was activated. This process was mediated by IL-6 secretion from the senescent osteoblasts which can promote osteoclast formation and bone resorption in the local microenvironment. IL-6 inhibition prevented osteoclast formation and reduced frequency of breast metastases. It is suggested that osteoclast mediated bone resorption provides a greater surface area for breast cancer cells to disseminate to [92]. Therapy-induced senescent cells located at distal sites to the tumor could therefore promote local niches for cancer cells to metastasize to. Senescent thyroid cells have been proposed to induce invasion and metastasis of surrounding non-senescent thyroid cancer cells *in vivo*. Orthotopic co-implantation of *BRAFV600E* induced senescent thyrocytes and thyroid cancer cells resulted in increased lymphovascular invasion and metastases in lymph nodes. This process was mediated via a CXCL12/CXCR4 axis [104](Figure 1). These results suggest senescence induced cancer cells could promote aggressive features of cancer cells which did not respond to therapy.

Recent reports have also proposed that chemotherapy-induced senescent cancer cells acquire stemness-like properties. Doxorubicin treatment in Eμ-Myc transgenic mice results in B-cell lymphoma senescence, but also to the upregulation of various stemness genes. Some of these

senescent stem-like lymphoma cells were able to re-enter the cell cycle and showed increased aggressiveness, mainly dependent on canonical Wnt signaling [103].



**Figure 2. Side effects caused by senescent cells and SASP induced by cancer therapy.** Cancer therapy is effective to inhibit cancer progression but can induce premature cellular senescence in normal or cancer cells. The SASP of senescent cells contribute to many side effects, including cardiac dysfunction, fatigue, bone loss and physical decline of the treated patients.

## 2) Non-tumorigenic effects

Treatment-related toxicities of cancer patients represent a major clinical issue. In the short term, these toxicities might lead to treatment discontinuation and decrease efficacy of the intervention. In the long term, adverse reactions and excessive damage to the therapy can cause cancer relapse and additional morbidities, including secondary tumors. This is becoming particularly relevant because of the increase number of cancer survivors [105], which decades after a successful anti-cancer treatment are suffering from a multitude of pathologies that they develop at an accelerated pace [106]. Additionally, survivors of childhood cancers are much more likely to develop secondary cancers, spinal disorders and pulmonary diseases in their adulthood [107].

Chronic inflammation is a predominant factor that not only influences the outcome of cancer treatment and contributes to cancer progression and cancer relapse, but also promotes potential side effects. Factors such as IL1A, IL6, IL8, CCL2 and CXCL12 are overexpressed after anti-cancer therapy [108] and have been associated with fatigue, cardiovascular morbidity, physical function decline and appetite loss [109–111]. Strikingly, these factors are also the major components of the SASP [14], and *in vivo* studies have shown that senescent cells are a major contributor to aging and age-related disease and dysfunctions, many observed in cancer patients [16,91]. Considering that senescence and SASP are promoted by many anti-cancer interventions (Table 1), it is logical to speculate that senescent cells might mediate part of the short- and long-term adverse reactions to cancer interventions.

A recent study from our laboratory demonstrated a correlation between detrimental senescent cells/SASP and the healthspan of chemotherapy-treated mice. Removal of chemotherapy-induced systemic senescent cells reduced the burden of inflammatory SASP factors and myelosuppression, thus preventing physical decline, frailty and cardiac dysfunctions [29]. Additionally, another group showed that inhibition of p38MAPK pathway in paclitaxel-treated mice improved the trabecular bone volume and bone density [112]. This beneficial effect is probably also mediated by inhibition of the SASP, as paclitaxel treatment could induce systemic senescence [29] and the p38MAPK pathway promotes SASP development [12]. This hypothesis was confirmed by another study showing that therapy-induced senescence contributed to bone loss, and that clearance of senescent cells rescued bone homeostasis [113] (Figure 2). Total body  $\gamma$ -irradiation of mice also induces senescence and an upregulated SASP

in various tissues [48,50]. Irradiated mice also developed early onset of symptoms associated with frailty including decreased cognition, weight loss and reduced grip strength [49].

- **Senotherapy in cancer**

Targeted removal of senescent cells (senolysis) is an emerging strategy in cancer treatments, in particular in combination with more traditional anti-cancer interventions [114]. ABT-263 (Navitoclax) is a pan-BCL2 inhibitor which interferes with senescence-associated anti-apoptosis and lead senescent cells to elimination [44]. Moreover, ABT-263 was able to kill senescence-like ovarian (OV4453 and OV1946) and breast (MDA-MB-231) cancer cells, in culture and in xenografts, pre-treated with the PARP inhibitor olaparib [115]. The bromodomain and extra-terminal (BET) protein BRD4 is overexpressed in triple negative breast cancer (TNBC), which makes the BET inhibitor a potential therapeutic agent [116]. Another pan-BCL2 family inhibitor, obatoclax, showed synergetic effects with BET inhibitor to kill senescence-like TNBC cells [116].

Mammalian Target Of Rapamycin (mTOR) is an essential regulator of the SASP [43,117] and mTOR inhibitors have shown senolytic effects in senescent cancer cells. Both concurrent and sequential treatments with chemotherapy (docetaxel) and the mTOR inhibitor temsirolimus (CCI-779) showed advanced anti-tumor effects in prostate (PC3) and breast (MDA-MB-468) cancer cells and xenografts [118]. A recent report showed that an inhibitor of the DNA-replication kinase CDC7, XL413, selectively induced senescence in liver cancer cells with p53 mutations (Huh7 and MHCC97H), and further administration of the mTOR inhibitor AZD8055 boosted the suppressive effects of tumor growth [119]. Since interfering with the SASP not always achieves senolysis, the lethal effect of mTOR inhibitors seem more dependent on the senescent cell type.

The histone deacetylase inhibitor panobinostat was also shown to promote senolysis of chemotherapy-treated non-small cell lung cancer (NSCLC) and head and neck squamous cell carcinoma (HNSCC) cell lines (A549, H460; UMSCC47, FaDu [120]. More recently, cardiac glycosides, in particular digoxin, were also characterized as universal senolytics able to kill various types of therapy-induced (bleomycin, gemcitabine, doxorubicin, etoposide and palbociclib) senescent cancer cells both *in vitro* and *in vivo* [121]. Inhibition of the SASP (senostasis) might also suffice to improve the response to cancer therapies. Rapamycin dramatically reduces growth of prostate cancer cell xenografts in mice treated with the

chemotherapeutic mitoxantrone [43]. Similar results were also seen in breast cancer xenografts when mice were treated with both doxorubicin and the SASP inhibitor metformin [122].

Together with potentiating cancer cell elimination, senotherapies have the potential to improve the healthspan of cancer patients. In a pre-clinical mouse model, ABT-263 cleared doxorubicin-induced senescent cells in mice bearing MMTV-PyMT breast carcinomas and significantly delayed cancer metastasis and recurrence [29]. Rapamycin and metformin also increase healthspan in middle-aged wild-type mice [123,124]. Administration of SASP inhibitors may therefore alleviate non-tumorigenic side effects associated with senescence inducing chemotherapy.

- **Can a foe turn into a friend?**

Besides the aforementioned detrimental effects, some of the proinflammatory SASP factors also hold the potential to contribute to the immune clearance of cancer cells. Restoration of p53 functions in lymphomas, sarcomas, and liver carcinomas leads to the features of cellular senescence and tumor regression [125,126], which is mainly mediated by the chronic inflammation and innate immune system triggered by the senescent stromal cells [126]. Immune surveillance by CD4<sup>+</sup> T cells of pre-malignant senescent hepatocytes also prevents the development of hepatocellular carcinomas [127], indicating that senescence and its secretory phenotypes provide an extrinsic barrier to malignancy in the early stages of cancer progression. Inactivation of Stat3 in a Pten-null prostate cancer model reduces levels of immunosuppressive SASP factors without affecting the expression of SASP chemoattractants, thus favoring immune-mediated reduction of tumor size. Moreover, treatment with a JAK2 inhibitor targeting the Stat3 pathway in combination with docetaxel modulated the chemotherapy-induced SASP and triggered a stronger immune-mediated anti-tumor response [128]. Recently, it has been shown in a mouse model of pancreatic ductal adenocarcinoma that a combination of MEK and CDK4/6 inhibitors induces cellular senescence, with the SASP favoring vascularization, drug delivery to the tumor site, and sensitivity to immunotherapy [129].

- **Concluding Remarks and Future Directions**

Cancer therapies inflicting high levels of DNA damage are still the primary treatment option for many malignancies, but they can inevitably lead to senescence induction and secretion of pro-inflammatory SASP molecules in both malignant and non-malignant cells. Using various *in vivo* models which genetically mimic human tumors, these SASP factors have been shown




to promote various hallmarks of tumorigenesis including proliferation, invasion, immunosuppression and metastasis. However, these studies are limited to few cancer types, and whether the findings are representative for most tumors needs to be addressed. This is particularly important because of the heterogeneous phenotypes of different senescence programs, for example the dependence of SASP composition on the cell type [130,131].

Inhibiting SASP production via the use of senolytics or senomorphics have been shown to suppress tumorigenesis and suppress the immunosuppressive environment promoted by senescent cancer cells. Moreover, senolysis might prevent the risk of some (pseudo)-senescent cancer cell to escape the growth arrest and resume uncontrolled proliferation. Thus, a ‘one-two punch’ strategy based on sequential pro- and anti-senescence interventions might be an effective strategy to treat cancer [119]. However, this approach requires more characterization of the response of senescent cancer cells to the senotherapy. For example, BCL-2 inhibitors ABT-263 and ABT-737 only results in a response in senescent cells that upregulate Bcl-xl or Bcl-w [6,7]. If a specific type of senescent cancer cell does not rely on these proteins for survival, they may not be effectively cleared via BCL-2 inhibitors.

Strategies towards senescence inducing therapies in old and young human cancer patients should be considered carefully (Figure 3). Young adults who have a low baseline of senescence prior to cancer treatment might benefit from adjuvant senotherapy that could reduce the likelihood of tumor relapse and premature aging phenotypes (Figure 3).

By contrast, older patients, which normally have a higher level of senescence pre-cancer treatment [132,133], might benefit from additional neoadjuvant senotherapy. Reducing the burst of senescence could potentially improve healthspan and prepare the patients to better cope with highly damaging stress deriving from the anti-cancer interventions. Senolytics or senomorphics would then be given again after the end of the anti-cancer therapy, to prevent further tumorigenesis from pro-inflammatory SASP factors (Figure 3). The feasibility of this approach should be tested further using appropriate cancer models of aged mice.



**Figure 3. Potential strategies for senotherapy mediated cancer treatment.** (A) Cancer therapy in young tumor microenvironment leads to cancer cell death and therapy-induced senescence (TIS) in cancer cells and stromal normal cells. The subsequent round of senotherapy selectively kills the senescent cancer and normal cells induced by therapy and a better treatment outcome is achieved after regeneration (green arrow route). (B) Senescent stromal normal cells massively exist in the old tumor microenvironment and contribute to cancer progression during the therapy and cancer relapse post the therapy. A second round of senotherapy is not able to achieve desired outcome as good as in young microenvironment (black arrow route). (C) An extra round of senotherapy before cancer therapy in the old tumor microenvironment to remove pre-exist senescent stromal normal cells will be beneficial. The rejuvenation step bridges the aged microenvironment to the young and renders a better treatment outcome (blue arrow route – green arrow route).

- **Acknowledgements:**

We thank the contributors from Science Illustration (<https://www.somersault1824.com/>) to share the components that we used to finish our figures. We are grateful to all the members of the Demaria lab for fruitful discussions.

- **References:**

- 1 Gorgoulis, V. *et al.* (2019) Cellular Senescence: Defining a Path Forward. *Cell* 179, 813–827
- 2 Hernandez-Segura, A. *et al.* (2018) Hallmarks of Cellular Senescence. *Trends Cell Biol* 28, 436–453
- 3 Dimri, G. and Medrano, E. (1995) A biomarker that identifies senescent human cells in culture and in aging skin in vivo. *Proc National Acad Sci* 92, 9363–9367
- 4 Hall, B. *et al.* (2017) p16(Ink4a) and senescence-associated  $\beta$ -galactosidase can be induced in macrophages as part of a reversible response to physiological stimuli. *Aging* 9, 1867–1884
- 5 Sharpless, N. and Sherr, C. (2015) Forging a signature of in vivo senescence. *Nat Rev Cancer* 15, 397–408
- 6 Yosef, R. *et al.* (2016) Directed elimination of senescent cells by inhibition of BCL-W and BCL-XL. *Nat Commun* 7, 11190
- 7 Zhu, Y. *et al.* (2016) Identification of a novel senolytic agent, navitoclax, targeting the Bcl-2 family of anti-apoptotic factors. *Aging Cell* 15, 428–435
- 8 Rodier, F. *et al.* (2010) DNA-SCARS: distinct nuclear structures that sustain damage-induced senescence growth arrest and inflammatory cytokine secretion. *J Cell Sci* 124, 68–81
- 9 Zhang, R. *et al.* (2007) Molecular Dissection of Formation of Senescence-Associated Heterochromatin Foci  $\nabla$   $\dagger$ . *Mol Cell Biol* 27, 2343–2358
- 10 Rodier, F. *et al.* (2009) Persistent DNA damage signalling triggers senescence-associated inflammatory cytokine secretion. *Nat Cell Biol* 11, 973–979
- 11 Chien, Y. *et al.* (2011) Control of the senescence-associated secretory phenotype by NF- $\kappa$ B promotes senescence and enhances chemosensitivity. *Gene Dev* 25, 2125–2136
- 12 Freund, A. *et al.* (2011) p38MAPK is a novel DNA damage response-independent regulator of the senescence-associated secretory phenotype. *Embo J* 30, 1536–1548
- 13 Kuilman, T. *et al.* (2008) Oncogene-Induced Senescence Relayed by an Interleukin-Dependent Inflammatory Network. *Cell* 133, 1019–1031
- 14 Coppé, J. *et al.* (2008) Senescence-Associated Secretory Phenotypes Reveal Cell-Nonautonomous Functions of Oncogenic RAS and the p53 Tumor Suppressor. *Plos Biol* 6, e301
- 15 Collado, M. *et al.* (2005) Tumour biology: senescence in premalignant tumours. *Nature* 436, 642
- 16 Poole, Rh. *et al.* (2002) DNA damage is able to induce senescence in tumor cells in vitro and in vivo. *Cancer Res* 62, 1876–83
- 17 Ewald, J. *et al.* (2010) Therapy-Induced Senescence in Cancer. *Jnci J National Cancer Inst* 102, 1536–1546

- 18 Campisi, J. *et al.* (2011) Cellular senescence: A link between cancer and age-related degenerative disease? *Semin Cancer Biol* 21, 354–359
- 19 Chabner, B. and Roberts, T. (2005) Timeline: Chemotherapy and the war on cancer. *Nat Rev Cancer* 5, 65–72
- 20 Wong, S. *et al.* (1998) Evidence of cisplatin-induced senescent-like growth arrest in nasopharyngeal carcinoma cells. *Cancer Res* 58, 5019–22
- 21 Yang, M. *et al.* (2012) Induction of Cellular Senescence by Doxorubicin Is Associated with Upregulated miR-375 and Induction of Autophagy in K562 Cells. *Plos One* 7, e37205
- 22 Roberson, R. *et al.* (2005) Escape from Therapy-Induced Accelerated Cellular Senescence in p53-Null Lung Cancer Cells and in Human Lung Cancers. *Cancer Res* 65, 2795–2803
- 23 Ota, H. *et al.* (2009) Sirolimus and Everolimus Induce Endothelial Cellular Senescence Via Sirtuin 1 Down-Regulation Therapeutic Implication of Cilostazol After Drug-Eluting Stent Implantation. *J Am Coll Cardiol* 53, 2298–2305
- 24 Jaiswal, A. *et al.* (2015) NSC666715 and Its Analogs Inhibit Strand-Displacement Activity of DNA Polymerase  $\beta$  and Potentiate Temozolomide-Induced DNA Damage, Senescence and Apoptosis in Colorectal Cancer Cells. *Plos One* 10, e0123808
- 25 Thorn, C. *et al.* (2011) Doxorubicin pathways: pharmacodynamics and adverse effects. *Pharmacogenet Genom* 21, 440–6
- 26 Calman, K. *et al.* (1980) Basic Principles of Cancer Chemotherapy. DOI: 10.1007/978-1-349-86135-4\_6
- 27 Mikula-Pietrasik, J. *et al.* (2019) Comprehensive review on how platinum- and taxane-based chemotherapy of ovarian cancer affects biology of normal cells. *Cell Mol Life Sci* 76, 681–697
- 28 Hu, W. *et al.* (2016) Mechanistic Investigation of Bone Marrow Suppression Associated with Palbociclib and its Differentiation from Cytotoxic Chemotherapies. *Clin Cancer Res* 22, 2000–2008
- 29 Demaria, M. *et al.* (2017) Cellular Senescence Promotes Adverse Effects of Chemotherapy and Cancer Relapse. *Cancer Discov* 7, 165–176
- 30 Scheule, R. and Perkins, R. (1992) Bleomycin stimulation of cytokine secretion by the human alveolar macrophage. *Am J Physiology* 262, L386-91
- 31 Muñoz-Espín, D. *et al.* (2018) A versatile drug delivery system targeting senescent cells. *Embo Mol Med* 10,
- 32 Qiu, T. *et al.* (2019) PTEN loss regulates alveolar epithelial cell senescence in pulmonary fibrosis depending on Akt activation. *Aging* 11,
- 33 Shivshankar, P. *et al.* (2012) Caveolin-1 Deficiency Protects from Pulmonary Fibrosis by Modulating Epithelial Cell Senescence in Mice. *Am J Resp Cell Mol* 47, 28–36
- 34 Aoshiba, K. *et al.* (2013) Senescence-associated secretory phenotype in a mouse model of bleomycin-induced lung injury. *Exp Toxicol Pathol* 65, 1053–1062
- 35 Comis, R. (1992) Bleomycin pulmonary toxicity: current status and future directions. *Semin Oncol* 19, 64–70
- 36 Baskar, R. *et al.* (2012) Cancer and Radiation Therapy: Current Advances and Future Directions. *Int J Med Sci* 9, 193–199

- 37 Chang, B. *et al.* (1999) A senescence-like phenotype distinguishes tumor cells that undergo terminal proliferation arrest after exposure to anticancer agents. *Cancer Res* 59, 3761–7
- 38 Zhang, X. *et al.* (2019) Promotion of cellular senescence by THG-1/TSC22D4 knockout through activation of JUNB. *Biochem Biophys Res Commun* 522, 897–902
- 39 Jones, K. *et al.* (2009) p53-Dependent accelerated senescence induced by ionizing radiation in breast tumour cells. *Int J Radiat Biol* 81, 445–458
- 40 Quick, Q. and Gewirtz, D. (2006) An accelerated senescence response to radiation in wild-type p53 glioblastoma multiforme cells. *J Neurosurg* 105, 111–118
- 41 Igarashi, K. *et al.* (2007) Radiation-induced senescence-like phenotype in proliferating and plateau-phase vascular endothelial cells. *Exp Cell Res* 313, 3326–3336
- 42 Ji, J. *et al.* (2015) Ionizing irradiation inhibits keloid fibroblast cell proliferation and induces premature cellular senescence. *J Dermatology* 42, 56–63
- 43 Laberge, R. *et al.* (2015) MTOR regulates the pro-tumorigenic senescence-associated secretory phenotype by promoting IL1A translation. *Nat Cell Biol* 17, 1049–1061
- 44 Chang, J. *et al.* (2016) Clearance of senescent cells by ABT263 rejuvenates aged hematopoietic stem cells in mice. *Nat Med* 22, 78–83
- 45 Zhu, Y. *et al.* (2015) The Achilles' heel of senescent cells: from transcriptome to senolytic drugs. *Aging Cell* 14, 644–658
- 46 Zhao, Y. *et al.* (2018) Naked mole rats can undergo developmental, oncogene-induced and DNA damage-induced cellular senescence. *Proc National Acad Sci* 115, 201721160
- 47 Chiche, A. *et al.* (2017) Injury-Induced Senescence Enables In Vivo Reprogramming in Skeletal Muscle. *Cell Stem Cell* 20, 407–414.e4
- 48 Demaria, M. *et al.* (2014) An Essential Role for Senescent Cells in Optimal Wound Healing through Secretion of PDGF-AA. *Dev Cell* 31, 722–733
- 49 Fielder, E. *et al.* (2019) Sublethal whole-body irradiation causes progressive premature frailty in mice. *Mech Ageing Dev* 180, 63–69
- 50 Le, O. *et al.* (2010) Ionizing radiation-induced long-term expression of senescence markers in mice is independent of p53 and immune status. *Aging Cell* 9, 398–409
- 51 Malumbres, M. and Barbacid, M. (2009) Cell cycle, CDKs and cancer: a changing paradigm. *Nat Rev Cancer* 9, 153–166
- 52 Yu, Q. *et al.* (2001) Specific protection against breast cancers by cyclin D1 ablation. *Nature* 411, 1017–1021
- 53 Yu, Q. *et al.* (2006) Requirement for CDK4 kinase function in breast cancer. *Cancer Cell* 9, 23–32
- 54 Rader, J. *et al.* (2013) Dual CDK4/CDK6 Inhibition Induces Cell-Cycle Arrest and Senescence in Neuroblastoma. *Am Assoc Cancer Res* 19, 6173–6182
- 55 Fry, D. *et al.* (2004) Specific inhibition of cyclin-dependent kinase 4/6 by PD 0332991 and associated antitumor activity in human tumor xenografts. *Mol Cancer Ther* 3, 1427–38
- 56 O'Leary, B. *et al.* (2016) Treating cancer with selective CDK4/6 inhibitors. *Nat Rev Clin Oncol* 13, 417–430

- 57 Coppé, J. *et al.* (2011) Tumor suppressor and aging biomarker p16(INK4a) induces cellular senescence without the associated inflammatory secretory phenotype. *J Biological Chem* 286, 36396–403
- 58 Guenther, L. *et al.* (2018) A Combination CDK4/6 and IGF1R Inhibitor Strategy for Ewing Sarcoma. *Clin Cancer Res Official J Am Assoc Cancer Res* 25, 1343–1357
- 59 Goel, S. *et al.* (2016) Overcoming Therapeutic Resistance in HER2-Positive Breast Cancers with CDK4/6 Inhibitors. *Cancer Cell* 29, 255–269
- 60 Goel, S. *et al.* (2017) CDK4/6 inhibition triggers anti-tumour immunity. *Nature* 548, 471–475
- 61 Gong, X. *et al.* (2017) Genomic Aberrations that Activate D-type Cyclins Are Associated with Enhanced Sensitivity to the CDK4 and CDK6 Inhibitor Abemaciclib. *Cancer Cell* 32, 761–776.e6
- 62 Torres-Guzmán, R. *et al.* (2014) Preclinical characterization of abemaciclib in hormone receptor positive breast cancer. *Oncotarget* 5, 69493–69507
- 63 Llanos, S. *et al.* (2019) Lysosomal trapping of palbociclib and its functional implications. *Oncogene* 38, 3886–3902
- 64 Guan, X. *et al.* (2017) Stromal Senescence By Prolonged CDK4/6 Inhibition Potentiates Tumor Growth. *Mol Cancer Res* 15, 237–249
- 65 Hari, P. *et al.* (2019) The innate immune sensor Toll-like receptor 2 controls the senescence-associated secretory phenotype. *Sci Adv* 5, eaaw0254
- 66 Dai, C. and Enders, G. (2000) p16INK4a can initiate an autonomous senescence program. *Oncogene* 19, 1613–1622
- 67 Patnaik, A. *et al.* (2016) Efficacy and Safety of Abemaciclib, an Inhibitor of CDK4 and CDK6, for Patients with Breast Cancer, Non–Small Cell Lung Cancer, and Other Solid Tumors. *Cancer Discov* 6, 740–753
- 68 Yoshida, A. *et al.* (2016) Induction of Therapeutic Senescence in Vemurafenib-Resistant Melanoma by Extended Inhibition of CDK4/6. *Cancer Res* 76, 2990–3002
- 69 Widodo, N. *et al.* (2007) Stress Chaperones, Mortalin, and Pex19p Mediate 5-Aza-2' Deoxycytidine-Induced Senescence of Cancer Cells by DNA Methylation-Independent Pathway. *Journals Gerontology Ser* 62, 246–255
- 70 Amatori, S. *et al.* (2011) Premature senescence induced by DNA demethylating agent (Decitabine) as therapeutic option for malignant pleural mesothelioma. *Lung Cancer* 71, 113–115
- 71 Putri, J. *et al.* (2017) Induction of senescence in cancer cells by 5'-Aza-2'-deoxycytidine: Bioinformatics and experimental insights to its targets. *Comput Biol Chem* 70, 49–55
- 72 Almeida, L. *et al.* (2017) Unlocking the chromatin of adenoid cystic carcinomas using HDAC inhibitors sensitize cancer stem cells to cisplatin and induces tumor senescence. *Stem Cell Res* 21, 94–105
- 73 Xu, W. *et al.* (2005) Induction of Polyploidy by Histone Deacetylase Inhibitor: A Pathway for Antitumor Effects. *Cancer Res* 65, 7832–7839
- 74 Elknerova, K. *et al.* (2011) Epigenetic modulation of gene expression of human leukemia cell lines - induction of cell death and senescence. *Neoplasma* 58, 35–44
- 75 Kaletsch, A. *et al.* (2018) Effects of novel HDAC inhibitors on urothelial carcinoma cells. *Clin Epigenetics* 10, 100

- 76 Chiao, M. *et al.* (2013) Suberoylanilide hydroxamic acid (SAHA) causes tumor growth slowdown and triggers autophagy in glioblastoma stem cells. *Autophagy* 9, 1509–26
- 77 Ogryzko, V. *et al.* (1996) Human fibroblast commitment to a senescence-like state in response to histone deacetylase inhibitors is cell cycle dependent. *Mol Cell Biol* 16, 5210–5218
- 78 Munro, J. *et al.* (2004) Histone deacetylase inhibitors induce a senescence-like state in human cells by a p16-dependent mechanism that is independent of a mitotic clock. *Exp Cell Res* 295, 525–538
- 79 Däbritz, J. *et al.* (2016) CD20-Targeting Immunotherapy Promotes Cellular Senescence in B-Cell Lymphoma. *Mol Cancer Ther* 15, 1074–1081
- 80 Krtolica, A. *et al.* (2001) Senescent fibroblasts promote epithelial cell growth and tumorigenesis: A link between cancer and aging. *Proc National Acad Sci* 98, 12072–12077
- 81 Fisher, D. *et al.* (2014) The two faces of IL-6 in the tumor microenvironment. *Semin Immunol* 26, 38–47
- 82 Kessenbrock, K. *et al.* (2010) Matrix Metalloproteinases: Regulators of the Tumor Microenvironment. *Cell* 141, 52–67
- 83 Waugh, D. and Wilson, C. (2008) The Interleukin-8 Pathway in Cancer. *Clin Cancer Res* 14, 6735–6741
- 84 Liu, D. and Hornsby, P. (2007) Senescent Human Fibroblasts Increase the Early Growth of Xenograft Tumors via Matrix Metalloproteinase Secretion. *Cancer Res* 67, 3117–3126
- 85 Canino, C. *et al.* (2012) SASP mediates chemoresistance and tumor-initiating-activity of mesothelioma cells. *Oncogene* 31, 3148–3163
- 86 MIAO, J. *et al.* (2014) Interleukin-6-induced epithelial-mesenchymal transition through signal transducer and activator of transcription 3 in human cervical carcinoma. *Int J Oncol* 45, 165–176
- 87 Goulet, C. *et al.* (2019) Cancer-associated fibroblasts induce epithelial–mesenchymal transition of bladder cancer cells through paracrine IL-6 signalling. *Bmc Cancer* 19, 137
- 88 Wang, L. *et al.* (2015) Activation of IL-8 via PI3K/Akt-dependent pathway is involved in leptin-mediated epithelial-mesenchymal transition in human breast cancer cells. *Cancer Biol Ther* 16, 1220–1230
- 89 Carmeliet, P. (2005) VEGF as a Key Mediator of Angiogenesis in Cancer. *Oncology* 69, 4–10
- 90 Coppé, J. *et al.* (2006) Secretion of Vascular Endothelial Growth Factor by Primary Human Fibroblasts at Senescence. *J Biol Chem* 281, 29568–29574
- 91 Sanoff, H. *et al.* (2014) Effect of Cytotoxic Chemotherapy on Markers of Molecular Age in Patients With Breast Cancer. *Jnci J National Cancer Inst* 106, dju057
- 92 Luo, X. *et al.* (2016) Stromal-Initiated Changes in the Bone Promote Metastatic Niche Development. *Cell Reports* 14, 82–92
- 93 Eggert, T. *et al.* (2016) Distinct Functions of Senescence-Associated Immune Responses in Liver Tumor Surveillance and Tumor Progression. *Cancer Cell* 30, 533–547
- 94 Jackson, J. *et al.* (2012) p53-Mediated Senescence Impairs the Apoptotic Response to Chemotherapy and Clinical Outcome in Breast Cancer. *Cancer Cell* 21, 793–806
- 95 ZHU, F. *et al.* (2014) Eotaxin-1 promotes prostate cancer cell invasion via activation of the CCR3-ERK pathway and upregulation of MMP-3 expression. *Oncol Rep* 31, 2049–2054

- 96 Chen, C. *et al.* (2019) CXCL5 induces tumor angiogenesis via enhancing the expression of FOXD1 mediated by the AKT/NF- $\kappa$ B pathway in colorectal cancer. *Cell Death Dis* 10, 178
- 97 Zhao, J. *et al.* (2017) Tumor-derived CXCL5 promotes human colorectal cancer metastasis through activation of the ERK/Elk-1/Snail and AKT/GSK3 $\beta$ / $\beta$ -catenin pathways. *Mol Cancer* 16, 70
- 98 Aldinucci, D. and Colombatti, A. (2014) The Inflammatory Chemokine CCL5 and Cancer Progression. *Mediat Inflamm* 2014, 292376
- 99 Jamaspishvili, T. *et al.* (2018) Clinical implications of PTEN loss in prostate cancer. *Nat Rev Urol* 15, 222–234
- 100 Chen, Z. *et al.* (2005) Crucial role of p53-dependent cellular senescence in suppression of Pten-deficient tumorigenesis. *Nature* 436, 725–730
- 101 Mitri, D. *et al.* (2014) Tumour-infiltrating Gr-1+ myeloid cells antagonize senescence in cancer. *Nature* 515, 134–137
- 102 Ruhland, M. *et al.* (2016) Stromal senescence establishes an immunosuppressive microenvironment that drives tumorigenesis. *Nat Commun* 7, 11762
- 103 Milanovic, M. *et al.* (2018) Senescence-associated reprogramming promotes cancer stemness. *Nature* 553, 96–100
- 104 Kim, Y. *et al.* (2017) Senescent tumor cells lead the collective invasion in thyroid cancer. *Nat Commun* 8, 15208
- 105 Miller, K. *et al.* (2019) Cancer treatment and survivorship statistics, 2019. *Ca Cancer J Clin* 69, 363–385
- 106 Zhu, J. *et al.* (2020) Accelerated aging in breast cancer survivors and its association with mortality and cancer recurrence. *Breast Cancer Res Tr* 180, 449–459
- 107 Bhakta, N. *et al.* (2017) The cumulative burden of surviving childhood cancer: an initial report from the St Jude Lifetime Cohort Study (SJLIFE). *Lancet* 390, 2569–2582
- 108 Mantovani, A. *et al.* (2008) Cancer-related inflammation. *Nature* 454, 436–444
- 109 Bhakta, N. *et al.* (2016) Cumulative burden of cardiovascular morbidity in paediatric, adolescent, and young adult survivors of Hodgkin's lymphoma: an analysis from the St Jude Lifetime Cohort Study. *Lancet Oncol* 17, 1325–1334
- 110 Cupit-Link, M. *et al.* (2017) Biology of premature ageing in survivors of cancer. *Esmo Open* 2, e000250
- 111 Hanahan, D. and Weinberg, R. (2011) Hallmarks of Cancer: The Next Generation. *Cell* 144, 646–674
- 112 Murali, B. *et al.* (2018) Inhibition of the Stromal p38MAPK/MK2 Pathway Limits Breast Cancer Metastases and Chemotherapy-Induced Bone Loss. *Cancer Res* 78, 5618–5630
- 113 Yao, Z. *et al.* (2020) Therapy-Induced Senescence Drives Bone Loss. *Cancer Res* 80, 1171–1182
- 114 Sieben, C. *et al.* (2018) Two-Step Senescence-Focused Cancer Therapies. *Trends Cell Biol* 28, 723–737
- 115 Fleury, H. *et al.* (2019) Exploiting interconnected synthetic lethal interactions between PARP inhibition and cancer cell reversible senescence. *Nat Commun* 10, 2556



- 116 Gayle, S. *et al.* (2019) Targeting BCL-xL improves the efficacy of bromodomain and extra-terminal protein inhibitors in triple-negative breast cancer by eliciting the death of senescent cells. *J Biol Chem* 294, 875–886
- 117 Herranz, N. *et al.* (2015) mTOR regulates MAPKAPK2 translation to control the senescence-associated secretory phenotype. *Nat Cell Biol* 17, 1205–1217
- 118 Fung, A. *et al.* (2009) Concurrent and Sequential Administration of Chemotherapy and the Mammalian Target of Rapamycin Inhibitor Temsirolimus in Human Cancer Cells and Xenografts. *Clin Cancer Res* 15, 5389–5395
- 119 Wang, C. *et al.* (2019) Inducing and exploiting vulnerabilities for the treatment of liver cancer. *Nature* 574, 268–272
- 120 Samaraweera, L. *et al.* (2017) A Novel Indication for Panobinostat as a Senolytic Drug in NSCLC and HNSCC. *Sci Rep-uk* 7, 1900
- 121 Triana-Martínez, F. *et al.* (2019) Identification and characterization of Cardiac Glycosides as senolytic compounds. *Nat Commun* 10, 4731
- 122 Iliopoulos, D. *et al.* (2011) Metformin Decreases the Dose of Chemotherapy for Prolonging Tumor Remission in Mouse Xenografts Involving Multiple Cancer Cell Types. *Cancer Res* 71, 3196–3201
- 123 Martin-Montalvo, A. *et al.* (2013) Metformin improves healthspan and lifespan in mice. *Nat Commun* 4, 2192
- 124 Bitto, A. *et al.* (2016) Transient rapamycin treatment can increase lifespan and healthspan in middle-aged mice. *Elife* 5, e16351
- 125 Ventura, A. *et al.* (2007) Restoration of p53 function leads to tumour regression in vivo. *Nature* 445, 661–665
- 126 Xue, W. *et al.* (2007) Senescence and tumour clearance is triggered by p53 restoration in murine liver carcinomas. *Nature* 445, 656–660
- 127 Kang, T. *et al.* (2011) Senescence surveillance of pre-malignant hepatocytes limits liver cancer development. *Nature* 479, 547–551
- 128 Toso, A. *et al.* (2014) Enhancing Chemotherapy Efficacy in Pten-Deficient Prostate Tumors by Activating the Senescence-Associated Antitumor Immunity. *Cell Reports* 9, 75–89
- 129 Ruscetti, M. *et al.* (2020) Senescence-Induced Vascular Remodeling Creates Therapeutic Vulnerabilities in Pancreas Cancer. *Cell* 181, 424–441.e21
- 130 Hernandez-Segura, A. *et al.* (2017) Unmasking Transcriptional Heterogeneity in Senescent Cells. *Curr Biol* 27, 2652–2660.e4
- 131 Basisty, N. *et al.* (2020) A proteomic atlas of senescence-associated secretomes for aging biomarker development. *Plos Biol* 18, e3000599
- 132 Baker, D. *et al.* (2016) Naturally occurring p16(Ink4a)-positive cells shorten healthy lifespan. *Nature* 530, 184–9
- 133 Liu, Y. *et al.* (2009) Expression of p16(INK4a) in peripheral blood T-cells is a biomarker of human aging. *Aging Cell* 8, 439–48

- 134 Bielak-Zmijewska, A. *et al.* (2014) A comparison of replicative senescence and doxorubicin-induced premature senescence of vascular smooth muscle cells isolated from human aorta. *Biogerontology* 15, 47–64
- 135 Chen, F. *et al.* (2018) Targeting SPINK1 in the damaged tumour microenvironment alleviates therapeutic resistance. *Nat Commun* 9, 4315
- 136 Jiang, C. *et al.* (2017) Serpine 1 induces alveolar type II cell senescence through activating p53-p21-Rb pathway in fibrotic lung disease. *Aging Cell* 16, 1114–1124
- 137 Baar, M. *et al.* (2017) Targeted Apoptosis of Senescent Cells Restores Tissue Homeostasis in Response to Chemotoxicity and Aging. *Cell* 169, 132–147.e16
- 138 Leontieva, O. *et al.* (2010) Weak p53 permits senescence during cell cycle arrest. *Cell Cycle Georget Tex* 9, 4323–7
- 139 Han, Z. *et al.* (2002) Role of p21 in apoptosis and senescence of human colon cancer cells treated with camptothecin. *J Biological Chem* 277, 17154–60
- 140 Peiris-Pagès, M. *et al.* (2015) Chemotherapy induces the cancer-associated fibroblast phenotype, activating paracrine Hedgehog-Gli signalling in breast cancer cells. *Oncotarget* 6, 10728–10745
- 141 Linge, A. *et al.* (2007) Downregulation of caveolin-1 affects bleomycin-induced growth arrest and cellular senescence in A549 cells. *Int J Biochem Cell Biology* 39, 1964–1974
- 142 Tian, Y. *et al.* (2019) Loss of PTEN induces lung fibrosis via alveolar epithelial cell senescence depending on NF-κB activation. *Aging Cell* 18, e12858
- 143 Terzis, A. (2003) Temozolomide induces apoptosis and senescence in glioma cells cultured as multicellular spheroids. *Brit J Cancer* 88, 463–469
- 144 Hirose, Y. *et al.* (2005) Akt Activation Suppresses Chk2-Mediated, Methylating Agent-Induced G2 Arrest and Protects from Temozolomide-Induced Mitotic Catastrophe and Cellular Senescence. *Cancer Res* 65, 4861–4869
- 145 Ohba, S. *et al.* (2009) Inhibition of c-Jun N-terminal kinase enhances temozolomide-induced cytotoxicity in human glioma cells. *J Neuro-oncol* 95, 307–316
- 146 Knizhnik, A. *et al.* (2013) Survival and Death Strategies in Glioma Cells: Autophagy, Senescence and Apoptosis Triggered by a Single Type of Temozolomide-Induced DNA Damage. *Plos One* 8, e55665
- 147 Mhaidat, N. *et al.* (2007) Temozolomide induces senescence but not apoptosis in human melanoma cells. *Brit J Cancer* 97, 1225–1233
- 148 DeMasters, G. *et al.* (2004) Potentiation of cell killing by fractionated radiation and suppression of proliferative recovery in MCF-7 breast tumor cells by the Vitamin D3 analog EB 1089. *J Steroid Biochem Mol Biology* 92, 365–374
- 149 Vijayaraghavan, S. *et al.* (2017) CDK4/6 and autophagy inhibitors synergistically induce senescence in Rb positive cytoplasmic cyclin E negative cancers. *Nat Commun* 8, 15916
- 150 Leontieva, O. and Blagosklonny, M. (2013) CDK4/6-inhibiting drug substitutes for p21 and p16 in senescence: Duration of cell cycle arrest and MTOR activity determine geroconversion. *Cell Cycle* 12, 3063–3069
- 151 Kovatcheva, M. *et al.* (2015) MDM2 turnover and expression of ATRX determine the choice between quiescence and senescence in response to CDK4 inhibition. *Oncotarget* 6, 8226–8243

- 152 Kovatcheva, M. *et al.* (2017) ATRX is a regulator of therapy induced senescence in human cells. *Nat Commun* 8, 386
- 153 Klein, M. *et al.* (2018) PDLIM7 and CDH18 regulate the turnover of MDM2 during CDK4/6 inhibitor therapy-induced senescence. *Oncogene* 37, 5066–5078
- 154 Valenzuela, C. *et al.* (2017) Palbociclib-induced autophagy and senescence in gastric cancer cells. *Exp Cell Res* 360, 390–396
- 155 Bollard, J. *et al.* (2017) Palbociclib (PD-0332991), a selective CDK4/6 inhibitor, restricts tumour growth in preclinical models of hepatocellular carcinoma. *Gut* 66, 1286
- 156 Schaer, D. *et al.* (2018) The CDK4/6 Inhibitor Abemaciclib Induces a T Cell Inflamed Tumor Microenvironment and Enhances the Efficacy of PD-L1 Checkpoint Blockade. *Cell Reports* 22, 2978–2994
- 157 Venturelli, S. *et al.* (2013) Differential Induction of Apoptosis and Senescence by the DNA Methyltransferase Inhibitors 5-Azacytidine and 5-Aza-2'-Deoxycytidine in Solid Tumor Cells. *Mol Cancer Ther* 12, 2226–2236
- 158 Borghesan, M. *et al.* (2016) DNA Hypomethylation and Histone Variant macroH2A1 Synergistically Attenuate Chemotherapy-Induced Senescence to Promote Hepatocellular Carcinoma Progression. *Cancer Res* 76, 594–606